

PCTWORLD INTELLECTUAL
Internat

INTERNATIONAL APPLICATION PUBLISHED U

WO 9602225A1

(51) International Patent Classification⁶ :

A61K 7/00, 7/50

A1

(43) International Publication Date: 1 February 1996 (01.02.96)

(21) International Application Number: PCT/GB95/01690

(22) International Filing Date: 18 July 1995 (18.07.95)

(30) Priority Data:
9414574.5 19 July 1994 (19.07.94) GB(71) Applicant (for AU BB CA GB IE KE LK MN MW NZ SD SG SZ
TT UG only): UNILEVER PLC [GB/GB]; Unilever House,
Blackfriars, London EC4P 4BQ (GB).(71) Applicant (for all designated States except AU BB CA GB IE
KE LK MN MW NZ SD SG SZ TT UG): UNILEVER N.V.
[NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).(74) Agents: LINN, S., Jonathan et al.; Mewburn Ellis, York House,
23 Kingsway, London WC2B 6HP (GB).(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH,
CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE,
KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN,
MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,
TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE,
CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT,
SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML,
MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ,
UG).**Published***With international search report.**Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.*

(54) Title: DETERGENT COMPOSITION

(57) Abstract

An aqueous liquid cleansing and moisturising composition comprising a surface active agent selected from anionic, nonionic, zwitterionic and cationic surface active agents, soap and mixtures thereof; a benefit agent having a weight average particle size in the range 50 to 500 microns; and a thickening agent. The thickening agent is added to the benefit agent in amount from 1 to 50 wt.%, based on the benefit agent.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

DETERGENT COMPOSITION

The present invention relates to detergent compositions suitable for care and personal washing of the skin. In particular, it relates to compositions which are formulated to give mild cleansing and conditioning of the skin.

Compositions formulated to cleanse the skin are well known. It is also known to formulate products which provide both a cleansing and a moisturising benefit.

For example WO 90/13283 discloses a composition comprising an acyl ester of an isethionic acid salt, a long chain fatty acid, a moisturiser component and, optionally, soap

One of the problems which may be encountered with such dual purpose compositions is that, whilst cleansing may be effective, there is an insufficient level of moisturising.

We have found a way of formulating such compositions such that they can deliver effective moisturising, conditioning or protection of the skin.

In WO 94/01085 and 94/01084 the advantage of depositing large particles of petrolatum from soap based compositions is recognised.

However, according to WO 94/03152, concerned with shower gels comprising, a non-soap detergent, silicone oil added to condition the skin and cationic polymers, the maximum average droplet size of the silicone oil that can be used is 2 microns, if product stability is to be maintained.

We have now found that larger particles, by particle is meant a solid particle or liquid droplet, of benefit agent can be incorporated into surface active containing

-2 -

compositions and effective delivery of the benefit agent can be achieved if the benefit agent is thickened.

Thus, according to the invention there is provided an aqueous liquid cleansing and moisturising composition comprising:-

- a) a surface active agent selected from anionic, nonionic, zwitterionic and cationic surface active agents, soap, and mixtures thereof;
- b) an benefit agent having a weight average particle size in the range 50 to 500 microns; and
- c) a thickening agent,

wherein the thickening agent is added to the benefit agent in amount from 1 to 50 wt% based on the benefit agent.

The composition is suitable for cleansing and "moisturising", "conditioning" or "protection" of the skin. The benefit agent is included in the composition to moisturise, condition and/or protect the skin. By "benefit agent" is meant a substance that softens the skin (stratum corneum) and keeps it soft by retarding the decrease of its water content and/or protects the skin.

Preferred benefit agents include

- a) silicone oils, gums and modifications thereof such as linear and cyclic polydimethylsiloxanes; amino, alkyl alkylaryl and aryl silicone oils;
- b) fats and oils including natural fats and oils such as jojoba, soyabean, rice bran, avocado, almond, olive, sesame, persic, castor, coconut, mink oils; cacao fat, beef tallow, lard; hardened oils obtained by hydrogenating the aforementioned

- 3 -

oils; and synthetic mono, di and triglycerides such as myristic acid glyceride and 2-ethylhexanoic acid glyceride;

- c) waxes such as carnauba, spermaceti, beeswax, lanolin and derivatives thereof;
- d) hydrophobic plant extracts;
- e) hydrocarbons such as liquid paraffins, petrolatum, microcrystalline wax, ceresin, squalene, squalane, and mineral oil;
- f) higher fatty acids such as lauric, myristic, palmitic, stearic, behenic, oleic, linoleic linolenic, lanolic, isostearic and polyunsaturated fatty acids (PUFA) acids;
- g) higher alcohols such as lauryl, cetyl, steryl, oleyl, behenyl, cholesterol and 2-hexadecanol alcohol;
- h) esters such as cetyl octanoate, myristyl lactate, cetyl lactate, isopropyl myristate, myristyl myristate, isopropyl palmitate, isopropyl adipate, butyl stearate, decyl oleate, cholesterol isostearate, glycerol monostearate, glycerol distearate, glycerol tristearate, alkyl lactate, for example lauryl lactate, alkyl citrate and alkyl tartrate;
- i) essential oils such as fish oils, mentha, jasmine, camphor, white cedar, bitter orange peel, ryu, turpentine, cinnamon, bergamont, citrus unshiu, calamus, pine, lavender, bay, clove, hiba, eucalyptus, lemon, starflower, thyme, peppermint, rose, sage, menthol, cineole, eugenol, citral, citronelle, borneol, linalool, geraniol, evening primrose, camphor, thymol, spirantol, pinene, limonene and terpenoid oils;
- j) lipids such as cholesterol, ceramides, sucrose esters and pseudo-ceramides as described in European Patent Specification No. 556 957;

- 4 -

- k) vitamins such as vitamin A and E, and vitamin alkyl esters, including those vitamin C alkyl esters;
- l) sunscreens such as octyl methoxyl cinnamate (Parsol MCX) and butyl methoxy benzoylmethane (Parsol 1789);
- m) Phospholipids; and
- n) mixtures of any of the foregoing components.

10 The benefit agent may be incorporated in a carrier in the compositions of the invention, particularly if it is likely to suffer detrimental interactions with other components of the composition. Benefit agents for which such detrimental interactions may occur include lipids; alkyl lactates;

15 sunscreens; esters such as isopropyl palmitate and isopropyl myristate; and vitamins. The carrier can, for example, be a silicone or hydrocarbon oil which is not solubilised/micellised by the surface active phase and in which the benefit agent is relatively soluble.

20 Particularly preferred benefit agents include silicone oils, gums and modifications thereof, esters such as isopropyl palmitate and myristate and alkyl lactates.

25 The benefit agent is preferably present in amount of from 0.1 to 15 wt%, most preferably from 0.2 to 10 wt%, more preferably from 0.5 to 7 wt%.

30 An advantage of the composition according to the invention is that, during use, it deposits benefit agent onto the skin at a level which results in a perceivable benefit. Without being bound by theory, it is believed the benefit agent is dispersed into large pools during dilution of the composition in use and these pools deposit readily onto the

35 skin.

- 5 -

In WO 94/01084 and WO 94/01085 the composition are structured by the presence of at least 5% insoluble solid fatty acid soap. However, it is believed that such high levels of solid phase material may adversely affect the amount of benefit agent deposited onto the skin. Thus, in the present invention it is preferred that the composition is substantially free of solid soap, by which is meant the level of solid soap is below 1wt%.

Suitable thickening agents for the benefit agent include polacrylates; fumed silica; natural and synthetic waxes, alkyl silicone waxes such as behenyl silicone wax; aluminium silicate; lanolin derivatives such as lanesterol; higher fatty alcohols; polyethylenecopolymers; narogel; polyammonium stearate; sucrose esters; hydrophobic clays; petrolatum; hydrotalcites; and mixtures thereof.

Hydrotalcites are materials of general formula



where M is a divalent metal ion e.g. Mg^{2+} ;
 N is a trivalent metal ion e.g. Al^{3+} ;
 X is an exchangeable anion e.g. CO_3^{2-} , NO_3^- ,
 stearate, cinnimate;
 m is the number of divalent metal ions; and
 n is the number of trivalent metal ions.

Particularly preferred thickening agents for the benefit agent include silica, alkyl silicone waxes, paraffin wax, higher fatty alcohols, petroleum jelly and polyethylenecopolymers.

Whilst some materials can function as both a benefit agent and a thickener therefor it will be appreciated that the benefit and thickening function cannot be provided by the

same component. However, it will be understood that where the composition comprises two or more benefit agents one of said benefit agents could also function as a thickening agent.

5

Preferably the amount of thickening agent is from 4 to 25% wt based on the level of benefit agent.

10

Although the compositions of the invention may be self-structuring preferably they will also comprise a structurant, i.e. a material added to increase the viscosity at zero shear. Suitable materials include swelling clays, for example laponite; fatty acids and derivatives thereof and, in particular fatty acid monoglyceride polyglycol ethers; cross-linked polyacrylates such as Carbopol (TM) (polymers available from Goodrich); acrylates and copolymers thereof, polyvinylpyrrolidone and copolymers thereof; polyethylene imines; salts such as sodium chloride and ammonium sulphate; sucrose esters; gellants; and mixtures thereof.

15

20

25

Of the clays particularly preferred are synthetic hectorite (laponite) clay used in conjunction with an electrolyte salt capable of causing the clay to thicken. Suitable electrolytes include alkali and alkaline earth salts such as halides, ammonium salts and sulphates.

30

35

The compositions according to the invention may also comprise a thickening agent in addition to the thickening agent added to the benefit agent, i.e. a material which maintains the viscosity of the composition as the shear rate thereof is increased during use. Suitable materials include cross-linked polyacrylates such as Carbopol (TM) (polymers available from Goodrich); fatty acids and derivatives thereof and, in particular, fatty acid monoglyceride polyglycol ethers; natural gums including


- 7 -

alginates, guar, xanthan and polysaccharide derivatives including carboxy methyl cellulose and hydroxypropyl guar; propylene glycols and propylene glycol oleates; salts such as sodium chloride and ammonium sulphate; glycerol tallowates; and mixtures thereof.

Further examples of structurants and thickeners are given in the International Cosmetic Ingredient Dictionary, Fifth Edition, 1993, published by CTFA (The Cosmetic, Toiletry & Fragrance Association), incorporated herein by reference.


The surface active agent can be selected from any known surfactant suitable for topical application to the human body. Mild surfactants, i.e. surfactants which do not damage the stratum corneum, the outer layer of skin, are particularly preferred.

One preferred anionic detergent is fatty acyl isethionate of formula:



where R is an alkyl or alkenyl group of 7 to 21 carbon atoms and M is a solubilising cation such as sodium, potassium, ammonium or substituted ammonium. Preferably at least three quarters of the RCO groups have 12 to 18 carbon atoms and may be derived from coconut, palm or a coconut/palm blend.

Another preferred anionic detergent is alkyl ether sulphate of formula:



where R is an alkyl group of 8 to 22 carbon atoms, n ranges from 0.5 to 10, especially from 1.5 to 8, and M is a solubilising cation as defined above.

Other possible anionic detergents include alkyl glyceryl ether sulphate, sulphosuccinates, taurates, sarcosinates, sulphoacetates, alkyl phosphate, alkyl phosphate esters and acyl lactylate, alkyl glutamates and mixtures thereof.

5

Sulphosuccinates may be monoalkyl sulphosuccinates having the formula: $R^5O_2CCH_2CH(SO_3M)CO_2M$; and amido-MEA sulphosuccinates of the formula:

10

$R^5CONHCH_2CH_2O_2CCH_2CH(SO_3M)CO_2M$; wherein R^5 ranges from C_8-C_{20} alkyl, preferably $C_{12}-C_{18}$ alkyl and M is a solubilising cation.

15

Sarcosinates are generally indicated by the formula: $R^5CON(CH_3)CH_2CO_2M$, wherein R^5 ranges from C_8-C_{20} alkyl, preferably $C_{12}-C_{18}$ alkyl and M is a solubilising cation.

20

Taurates are generally identified by the formula: $R^5CONR^6CH_2CH_2SO_3M$, wherein R^5 ranges from C_8-C_{20} alkyl, preferably $C_{12}-C_{18}$ alkyl, R^6 ranges from C_1-C_4 alkyl, and M is a solubilising cation.

Harsh surfactants such as primary alkane sulphonate or alkyl benzene sulphonate will generally be avoided.

25

30

Suitable nonionic surface active agents include alkyl polysaccharides, lactobionamides, ethyleneglycol esters, glycerol monoethers, polyhydroxyamides (glucamide), primary and secondary alcohol ethoxylates, especially the C_{8-20} aliphatic alcohols ethoxylated with an average of from 1 to 20 moles of ethylene oxide per mole of alcohol.

35


If the surface active agent comprises soap, the soap is preferably derived from materials with a C_8 to C_{22} substantially saturated carbon chain and, preferably, is a potassium soap with a C_{12} to C_{18} carbon chain.

- 9 -

Mixtures of any of the foregoing surface active agents may also be used.

The surface active agent is preferably present at a level of from 1 to 35 wt%, preferably 3 to 30 wt%.

It is also preferable that the composition includes from 0.5 to 15 wt% of a cosurfactant agent with skin-mildness benefits. Suitable materials are zwitterionic detergents which have an alkyl or alkenyl group of 7 to 18 carbon atoms and comply with an overall structural formula:-



where R¹ is alkyl or alkenyl of 7 to 18 carbon atoms;
R² and R³ are each independently alkyl, hydroxyalkyl or carboxyalkyl of 1 to 3 carbon atoms;


m is 2 to 4;

n is 0 or 1;

X is alkylene of 1 to 3 carbon atoms optionally substituted with hydroxyl, and


Y is -CO₂⁻ or -SO₃⁻.

Zwitterionic detergents within the above general formula include simple betaines of formula:-



and amido betaines of formula:

- 10 -



5


where m is 2 or 3.

In both formulae R^1 , R^2 and R^3 are as defined previously. R^1 may, in particular, be a mixture of C_{12} and C_{14} alkyl groups derived from coconut so that at least half, preferably, at least three quarter of the groups R^1 has 10 to 14 carbon atoms. R^2 and R^3 are preferably methyl.

10


A further possibility is a sulphobetaine of formula:-

15




20

or



25

where m is 2 or 3, or variants of these in which $-(\text{CH}_2)_n \text{SO}_3^-$ is replaced by



30

R^1 , R^2 and R^3 in these formulae are as defined previously.

Furthermore, the benefit agent may also function as a carrier to deliver efficacy agents to skin treated with the compositions of the invention. This route is particularly useful for delivering efficacy agents which are difficult to deposit onto the skin or those which suffer detrimental interactions with other components in the composition. In such cases the carrier is a often a silicone or hydrocarbon

35

oil which is not solubilised/micellised by the surface active phase and in which the efficacy agent is relatively soluble. Examples of such efficacy agents include anti-viral agents; hydroxycaprylic acids; pyrrolidone; carboxylic acids; 2,4,4'-trichloro-2'-hydroxydiphenyl ether (Irgasan DP300); 3,4,4'-trichlorocarbanilide; salicylic acid; benzoyl peroxide; perfumes; essential oils; germicides and insect repellants such as N,N-dimethyl m-toluamide (DEET); and mixtures thereof.

Compositions of the invention may be formulated as products for washing the skin, for example bath or shower gels, hand washing compositions, facial washing liquids; pre-and post-shaving products; rinse-off, wipe-off and leave-on skin-care products.

The compositions of the invention will generally be pourable liquids or semi-liquids eg pastes and will preferably have a viscosity in the range 1000 to 100,000 mPas measured at a shear rate 10 s^{-1} and 25°C in a Haake Rotoviscometer RV20.

The compositions of the invention will preferably exhibit a Newtonian viscosity at a shear stress of 0.01 Pa, at 25°C , of at least 5,000 Pas, preferably greater than 10,000 Pas.

The above-mentioned characteristic viscosity measurements may be determined exactly (as in the case of the non-zero shear viscosities) using, for example a Carrimed CSL 100 low stress rheometer or obtained from an extrapolation according to the Cross Model, (see J of the Chemical Engineer, 1993, paper entitled "Rheology for the Chemical Engineer" by H Barnes) as in the case of zero shear rate.

Other typical components of the compositions include opacifiers, preferably 0.2 to 2.0 wt%; preservatives,

preferably 0.2 to 2.0 wt% and perfumes, preferably 0.5 to 2.0 wt%.

5 According to a further aspect of the invention there is provided a process for preparing compositions according to the invention comprising:-

- 10 a) forming a base composition comprising at least one surface active agent selected from anionic, nonionic, zwitterionic and cationic surface active agents, soap and mixtures thereof;
- 15 b) adding a thickening agent to the benefit agent in an amount from 1 to 50%wt, based on the benefit agent; and
- c) mixing the base formulation of step a) with the benefit agent.

20 The invention will be further illustrated by reference to the following non-limiting examples.

Examples

- 25 In the examples:-
- Alkylpolyglucoside was Plantaren 2000 ex Henkel
- Behenyl alcohol was Nacol 22-97 ex Condea.
- Behenyl silicone wax was Cire 71649 ex Rhone Poulenc.
- Coco amidopropyl betaine was Amonyl BA 380 ex Seppic.
- 30 Cross-linked polyacrylate was Carbopol ETD 2020 ex Goodrich.
- Guar hydroxypropyl trimonium chloride was Jaguar C-13-S ex Meyhall.
- Hydrotalcite was hydrotalcite containing IPP, Multigel IPP
- 35 ex Guilini Chemie GmbH
- IPP (isopropyl palmitate) was ESTOL 1517 ex Unichema.

- 13 -

Lauryl lactate was Crodamol LL ex Croda Chemicals.

MEA suphosuccinate was witco 5690 ex Witco.

Polyethylene AC617 was from A-C Performance Additives.

Silica was a hydrophobically modified silica, Aerosil R972
ex Degussa.

The silicone oils were all DC200, a polydimethylsiloxane ex
Dow Corning, but with a varying viscosities as stated
below.

Silicone oil emulsion was BC 92/057 ex Basildon.

Sodium lauryl ether sulphate was Genapol ZRO ex Hoechst

Example I

In this example, the effect of thickening a silicone oil
benefit agent with silica on the deposition of silicone oil
onto the skin was examined.

The following method was used to determine the amount of
benefit agent deposited onto full thickness porcine skin (5
x 15 cm) treated with compositions according to the
invention.

The skin was prehydrated and then 0.5 ml of the product
applied to it. The product was lathered for 10 seconds and
then rinsed for 10 seconds under running water.

Thereafter the skin was wiped once with a paper towel to
remove excess water.

2 minutes after drying a strip of adhesive tape was pressed
onto the skin for 30 seconds by applying a constant load of
100g.cm⁻². The adhesive tape employed was J-Lar Superclear
(TM) tape having a width of 2.5cm. In total ten strips of
tape were applied to adjacent sites on the skin.

- 14 -

In this test procedure silicone which has deposited on the skin will subsequently be transferred to the tape along with some of the outer layer of the skin.

5 The amounts of silicon and skin adhering to the tape are determined by means of X-ray fluorescence spectroscopy. The tape strips are placed in an X-ray fluorescence spectrometer with the adhesive side facing the beam of this machine. A mask is applied over the tape to define a
10 standardised area in the middle of the tape which is exposed to the X-ray beam. The sample chamber of the machine is placed under vacuum before making measurements and the spectrometer is then used to measure the quantities of silicon and sulphur. The sulphur is representative of
15 the amount of skin which has transferred to the tape.

The amounts of silicon and sulphur observed with a clean piece of adhesive tape are subtracted from the experimental measurements. The experimental measurements for the
20 average levels of sulphur and silicon are expressed as a ratio of silicon to sulphur. From this ratio it is possible to determine silicone oil deposition per unit area of skin.

25 A base shower gel formulation having the following composition was prepared.

	<u>% wt</u>
Sodium lauryl ether sulphate	4.00
Sodium coco amido propyl betaine	1.00
30 Alkylpolyglucoside	5.00
Sorbic acid	0.37
Trisodium citrate dihydrate	0.49
Cross-linked polyacrylate	0.9
Thickener*	~1
35 5N NaOH**	~1
Water + minors	to 100

- 15 -

* Thickener added to give the required viscosity ($\sim 5000 \text{ mPas}$ at 10 s^{-1} and 25°C)

** NaOH was added to adjust the pH of the composition to pH 5.3.

5

10

The composition was prepared by dispersing the polyacrylate in excess water. Thereafter sorbic acid and trisodium citrate dihydrate were added to the resulting polymer dispersion. The three surface active agents were mixed and the resulting mix added to the polymer dispersion. Thereafter minors were added. The viscosity of the resulting composition was measured at a shear rate of 10 s^{-1} at 25°C and thickener added until the required viscosity obtained.

15

20

Two Harvard 44 syringe pumps were used to infuse the base formulation and the benefit agent, silicone oil. Silicone oil was present at a level of 5 wt% based on the total composition. One syringe was filled with the base formulation and the other with silicone oil. The syringes were then inserted into the syringe pump and the infusion rate set at 5:95 oil:base. The oil and base were forced through a static in-line mixer and a composition with oil particles of the required size obtained. The size of the particles can be controlled by the diameter of the static mixer, the flow rate and length of the mixer tube. The size of the particles can be determined using a Malvern Mastersizer.

25

30

Silicone deposition from the compositions according to the invention was measured by the test procedure described above and compared with that from a similar composition (A) comprising a silicone oil emulsion having a viscosity of 60000 mPas at 10 s^{-1} and 25°C .

35

Comparative Composition (Composition A)

		<u>%wt</u>
5	Sodium lauryl ether sulphate	13.00
	Coco amidopropyl betaine (CAPB)	2.00
	Silicone oil emulsion	5.00
	Guar hydroxypropyl trimonium chloride	0.10
10	Sorbic acid	0.37
	Sodium citrate dihydrate	0.49
	Sodium chloride'	~2
	Citric acid''	~0.01
	Water + minors	to 100

15

level can be varied in order to give the required viscosity of 5000 mPas at 10s⁻¹ and 25°C.

level can be varied in order to give the required pH.

20

The following results were obtained.

	<u>%Silica</u>	<u>Si:S</u>	<u>particle</u>
	<u>added</u>	<u>ratio</u>	<u>size/um</u>
Comparison	0	0.9	0.5
25 Ia	8.3	63.9	424
Ib	9.7	22.2	354
Ic	4.7	21.3	341
Id	6.1	11.5	289

30

The results demonstrate the advantage, in terms of deposition of silicone, using a composition comprising a thickened silicone oil over a composition comprising a unthickened silicone oil.

35

Example II

Example I was repeated except that the benefit agent used was IPP (isopropylpalmitate). The IPP was thickened with three different thickeners at a level of 10% by weight based on the IPP.

The following method was used to determine the amount of IPP deposited onto full thickness porcine skin (5 x 15cm pieces) treated with compositions according to the invention. The skin was treated and washed with the compositions according to the invention by the same method as described for example I. Thereafter the skin was extracted with ethanol three times (3 ml per extract) and the extract made up to 10g and submitted for Gas Chromatography analysis to determine the amount of IPP deposited.

The following results were obtained:-

<u>thickener</u>	<u>deposition/ppm</u>
none	6.1
polyethylene AC617	441.9
behenyl silicone wax	138.1
hydrotalcite	59.1

The results demonstrate the increased deposition of IPP obtained when the IPP is thickened.

Example III

In this example the benefit agent used was lauryl lactate.

The following method was used to determine the amount of lauryl lactate deposited onto skin treated with compositions according to the invention containing lauryl lactate.

- 18 -

Porcine full thickness skin was prehydrated and then 0.20g of the product and 0.8g of water applied to it. The skin was washed for 30 seconds and then rinsed for 10 seconds under running water.

5

Thereafter the skin was wiped once with a paper towel to remove excess water. 30 seconds after drying 5 strips of Desquame tape were applied to the skin sequentially for 10 seconds.

10

The strips were then removed from the skin and hydrolysed in 2ml of NaOH (0.5M) at 60°C for 60 minutes. They were then neutralised with 2ml HCl (0.5M) and in Sorensens phosphate buffer (pH 7.0). The total lauryl lactate and the protein content of the tapes, representative of the lauryl lactate transferred to the skin, was determined using Sigma diagnostic assay Kits 735/10 and 690. A solution containing the assay and treated strips was prepared and the absorbance at 540nm measured in a UV spectrometer.

15
20

The base formulation for the following experiments was:-

	<u>Wt%</u>
25 MEA sulphosuccinate	11.7
Cross-linked polyacrylate	0.3
Sodium cocoyl isethionate	8.5
Coco amidopropyl betaine	15.8
Water and minors	to 100

30

It was prepared by dispersing the polyacrylate in excess water. A premix of the three surfactants was then added to the polymer dispersion. Thereafter minors were added.

35

The following benefit agent/thickener combination was used:-

- 19 -

		<u>wt%</u>	<u>Droplet size/μm</u>
x	Lauryl lactate	85	172.2
	Polyethylene AC617	15	

The mixture X was prepared by heating the two components together to a temperature of -110°C to melt in the polyethylene. The mixture was then allowed to cool before being used.

An emulsion of the base formulation and thickened benefit agent, was applied to the skin using the test procedure described above.

The following results were obtained:-

		<u>lactate/Absorbance units ($\times 10^{-3}$)</u>
20	Untreated skin	81
	base formulation (no lauryl lactate)	63
	base + 5% X	128
	base + 5% lauryl lactate	102
	base + 10% X	154
25	base + 10% lauryl lactate	107
	* based on the total composition	

In a separate set of experiments, but with the same base formulation as described above, the effect of a range of thickening agents (all at a level of 5%wt based on the lauryl lactate) on the deposition of lauryl lactate was examined. The lauryl lactate or thickened lauryl lactate was added to the base formulation at a level of 5%wt.

- 20 -

		<u>lactate/Absorbance</u>
		<u>units (x10⁻³)</u>
	lauryl lactate (no thickener)	112
	lauryl lactate + paraffin wax	148
5	lauryl lactate + behenyl alcohol	152
	lauryl lactate + petroleum jelly	148
	lauryl lactate + polyethylene AC617	129
	lauryl lactate + behenyl silicone wax	167
10	The results demonstrate the benefit of thickening the benefit agent with a thickening agent	

CLAIMS

1. An aqueous liquid cleansing and moisturising composition comprising:-

- 5
- a) a surface active agent selected from anionic, nonionic, zwitterionic and cationic surface active agents, soap and mixtures thereof;
- 10
- b) an benefit agent having a weight average particle size in the range 50 to 500 microns; and
- c) a thickening agent.

15

wherein the thickening agent is added to the benefit agent in amount from 1 to 50%wt, based on the benefit agent.

2. A composition according to claim 1 wherein the thickening agent is selected from polyacrylates; silica, natural and synthetic waxes, alkyl silicone waxes; aluminium silicate; lanolin derivatives; higher fatty alcohols; polyethylenecopolymers; narogel; polyammonium stearate; sucrose esters; hydrophobic clays; petrolatum; hydrotalcites; and mixtures thereof.

25

3. A composition according to claim 1 wherein the composition is structured with a surfactant selected from swelling clays; fatty acid and derivatives thereof; cross-linked polyacrylates; acrylates and copolymers thereof; polyvinylpyrrolidone and copolymers thereof; polyethylene imines; salts; sucrose esters; gellants; and mixtures thereof.

30

4. A composition according to claim 1 wherein the benefit agent is selected from silicone oils; gums; fats; oils; waxes; hydrophobic plant extracts; hydrocarbons; fatty

35

acids; alcohols; esters; essential oils; lipids;
phospholipids; vitamins; sunscreens; and mixtures thereof.

5 5. A composition according to any one of the preceding
claims comprising 0.1 to 15 wt% of the benefit agent.

10 6. A composition according to any one of the preceding
claims wherein the benefit agent functions as a carrier to
deliver efficacy agents to skin treated with the
composition.

7. A composition according to any one of the preceding
claims comprising 1 to 35 wt% of the surface active agent.

15 8. A composition according to any one of the preceding
claims further comprising a cosurfactant selected from
betaines, amidobetaines and sulphobetaines.

20 9. A process for preparing a composition according to
claim 1 comprises:-

25 a) forming a base composition comprising at least one
surface active agent selected from anionic, nonionic,
zwitterionic and cationic surface active agents, soap and
mixtures thereof;

b) adding a thickening agent to the benefit agent in an
amount from 1 to 50 wt%, based on the benefit agent; and

30 c) mixing the base formulation of step a) with the
benefit agent.

35 10. A method of depositing a benefit agent from an aqueous
liquid cleansing and moisturising composition, the method
comprising providing the said benefit agent in a
composition comprising;

- 23 -

- a) a surface active agent selected from anionic, nonionic, zwitterionic and cationic surface active agents, soap and mixtures thereof;
- 5 b) an benefit agent having a weight average particle size in the range 50 to 500 microns; and
- c) a thickening agent, and wherein the thickening agent is added to the benefit agent in amount from 1 to 50% wt, based on the benefit agent.
- 10

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/GB 95/01690

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K7/00 A61K7/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,94 01084 (THE PROCTER & GAMBLE COMPANY) 20 January 1994 cited in the application see the whole document ---	1-10
X	EP,A,0 452 202 (L'OREAL) 16 October 1991 see the whole document ---	1-10
X	WO,A,93 19149 (THE PROCTER & GAMBLE COMPANY) 30 September 1993 see the whole document ---	1-10
P,X	WO,A,94 17166 (THE PROCTER & GAMBLE COMPANY) 4 August 1994 see the whole document ---	1-10
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "A" document member of the same patent family

Date of the actual completion of the international search

14 November 1995

Date of mailing of the international search report

24 11 95

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2240 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Couckuyt, P

INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/GB 95/01690

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,93 09761 (RICHARDSON-VICKS INC.) 27 May 1993 see the whole document ---	1-10
X	WO,A,93 21293 (THE PROCTER & GAMBLE COMPANY) 28 October 1993 see the whole document ---	1-10
X	EP,A,0 268 982 (TORAY SILICONE COMPANY LTD) 1 June 1988 see the whole document ---	1-10
X	EP,A,0 407 042 (COLGATE-PALMOLIVE COMPANY) 9 January 1991 see the whole document ---	1-10
X	EP,A,0 413 417 (COLGATE-PALMOLIVE COMPANY) 20 February 1991 see the whole document ---	1-10
X	EP,A,0 552 024 (UNILEVER PLC) 21 July 1993 see the whole document ---	1-10
A	EP,A,0 485 212 (UNILEVER PLC) 13 May 1992 see the whole document -----	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/GB 95/01690

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9401084	20-01-94	US-A- 5308526 JP-T- 7508752	03-05-94 28-09-95
EP-A-452202	16-10-91	FR-A- 2660554 AU-B- 630026 AU-B- 7408991 DE-D- 69100670 DE-T- 69100670 ES-T- 2060316 JP-A- 5345706	11-10-91 15-10-92 10-10-91 05-01-94 05-05-94 16-11-94 27-12-93
WO-A-9319149	30-09-93	AU-B- 3811193 CA-A- 2131174 CN-A- 1078746 EP-A- 0636166	21-10-93 30-09-93 24-11-93 01-02-95
WO-A-9417166	04-08-94	NONE	
WO-A-9309761	27-05-93	AU-A- 3136593 CA-A- 2122272 EP-A- 0613369 JP-T- 7501077 PT-A- 101082 US-A- 5439682	15-06-93 27-05-93 07-09-94 02-02-95 28-02-94 08-08-95
WO-A-9321293	28-10-93	NONE	
EP-A-268982	01-06-88	JP-B- 4062288 JP-A- 63130512 AU-B- 615423 AU-B- 8327187 DE-A- 3777763 WO-A- 8803792	05-10-92 02-06-88 03-10-91 16-06-88 30-04-92 02-06-88
EP-A-407042	09-01-91	US-A- 5051250 US-A- 4997641 US-A- 5213716 AT-T- 118165 AT-T- 125689 AU-B- 5755890	24-09-91 05-03-91 25-05-93 15-02-95 15-08-95 03-01-91

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 95/01690

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-407042		CA-A- 2019341	21-12-90
		CA-A- 2019352	21-12-90
		CA-A- 2019358	21-12-90
		CN-A- 1048422	09-01-91
		DE-D- 69016715	23-03-95
		DE-T- 69016715	28-09-95
		DE-D- 69021288	07-09-95
		EP-A- 0407040	09-01-91
		EP-A- 0407041	09-01-91
		EP-A- 0413416	20-02-91
		EP-A- 0413417	20-02-91
		GR-A- 90100471	15-11-91
		JP-A- 3051367	05-03-91
		US-A- 5348736	20-09-94
		US-A- 5415857	16-05-95
		US-A- 5346642	13-09-94
		AU-B- 640382	26-08-93
		AU-B- 5768890	16-05-91
		CA-A- 2019346	21-12-90
		CN-A- 1051501	22-05-91
		GR-B- 1000728	23-11-92
		HU-B- 210429	28-04-95
		JP-A- 3153619	01-07-91
		NZ-A- 234189	25-11-94
		PL-B- 165297	30-12-94
		TR-A- 26442	15-03-95
		AU-B- 635749	01-04-93
		AU-B- 5755990	16-05-91
		CN-A- 1051599	22-05-91
		GR-B- 1000696	08-10-92
		HU-B- 210123	28-02-95
		JP-A- 3153620	01-07-91
		NZ-A- 234190	26-08-94
		TR-A- 26421	15-03-95
		AU-B- 642651	28-10-93
		AU-B- 5753390	10-10-91
		CN-A- 1055481	23-10-91
		GR-B- 1000710	08-10-92
		HU-B- 209692	28-10-94
		JP-A- 3291213	20-12-91

INTERNATIONAL SEARCH REPORT
Information on patent family members

International Application No
PCT/GB 95/01690

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-407042		NO-B- 177923	11-09-95
		NZ-A- 234191	26-08-94
		PL-B- 164567	31-08-94
		TR-A- 26429	15-03-95
<hr/>			
EP-A-413417	20-02-91	US-A- 5051250	24-09-91
		US-A- 4997641	05-03-91
		US-A- 5213716	25-05-93
		AT-T- 118165	15-02-95
		AT-T- 125689	15-08-95
		AU-B- 5755890	03-01-91
		CA-A- 2019341	21-12-90
		CA-A- 2019352	21-12-90
		CA-A- 2019358	21-12-90
		CN-A- 1048422	09-01-91
		DE-D- 69016715	23-03-95
		DE-T- 69016715	28-09-95
		DE-D- 69021288	07-09-95
		EP-A- 0407040	09-01-91
		EP-A- 0407041	09-01-91
		EP-A- 0413416	20-02-91
		EP-A- 0407042	09-01-91
		GR-A- 90100471	15-11-91
		JP-A- 3051367	05-03-91
		US-A- 5348736	20-09-94
		US-A- 5415857	16-05-95
		US-A- 5346642	13-09-94
		AU-B- 640382	26-08-93
		AU-B- 5768890	16-05-91
		CA-A- 2019346	21-12-90
		CN-A- 1051501	22-05-91
		GR-B- 1000728	23-11-92
		HU-B- 210429	28-04-95
		JP-A- 3153619	01-07-91
		NZ-A- 234189	25-11-94
		PL-B- 165297	30-12-94
		TR-A- 26442	15-03-95
		AU-B- 635749	01-04-93
		AU-B- 5755990	16-05-91
		CN-A- 1051599	22-05-91

INTERNATIONAL SEARCH REPORT
Information on patent family members

Int ional Application No
PCT/GB 95/01690

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-413417		GR-B- 1000696	08-10-92
		HU-B- 210123	28-02-95
		JP-A- 3153620	01-07-91
		NZ-A- 234190	26-08-94
		TR-A- 26421	15-03-95
		AU-B- 642651	28-10-93
		AU-B- 5753390	10-10-91
		CN-A- 1055481	23-10-91
		GR-B- 1000710	08-10-92
		HU-B- 209692	28-10-94
		JP-A- 3291213	20-12-91
		NO-B- 177923	11-09-95
		NZ-A- 234191	26-08-94
		PL-B- 164567	31-08-94
		TR-A- 26429	15-03-95
EP-A-552024	21-07-93	AU-A- 3181393	22-07-93
		CA-A- 2087140	16-07-93
		JP-A- 5279232	26-10-93
EP-A-485212	13-05-92	AT-T- 125150	15-08-95
		AU-B- 654154	27-10-94
		AU-B- 8696791	14-05-92
		AU-B- 644031	02-12-93
		AU-B- 8861991	11-06-92
		DE-D- 69111383	24-08-95
		EP-A- 0509079	21-10-92
		WO-A- 9208440	29-05-92
		JP-A- 4283509	08-10-92
		JP-T- 5503312	03-06-93
		ZA-A- 9108840	07-05-93

THIS PAGE BLANK (USPTO)